arXiv:1512.03528v1 [math.AT] 11 Dec 2015

Circle actions with two fixed points

Oleg R. Musin*

Abstract

We prove that if the circle group acts smooth and unitary on $2n$–dimensional stably complex manifold with two isolated fixed points and it is not bound equivariantly, then $n = 1$ or 3. Our proof relies on the rigid Hirzebruch genera.

1 Introduction

We say that M is a *unitary* $\mathbf{S}^1$*–manifold* if it is a smooth closed manifold with an effective circle action such that its tangent bundle admits an $\mathbf{S}^1$–equivariant stable complex structure.

Let M be a unitary $2n$–dimensional $\mathbf{S}^1$–manifold with m isolated fixed points $p_1, \dots, p_m$. Denote weights in p_i by $w_{i1}, \dots, w_{in}$ and its sign by ε_i, where $\varepsilon_i = \pm 1$.

If $m = 2$, then known the following sets of weights for unitary $\mathbf{S}^1$–manifolds:

(Z): $w_{1i} = w_{2i} = a_i$, where $a_i \in \mathbb{Z}$, $a_i \neq 0$, for all $i = 1, \dots, n$, and $\varepsilon_1 = -\varepsilon_2$. A manifold with these weights is cobordant to zero in the ring $U_*^{\mathbf{S}^1}$ of complex bordisms with circle actions.

(L_1): $n = 1$, $w_{11} = a$, $w_{21} = -a$, where $a > 0$, and $\varepsilon_1 = \varepsilon_2$. It is a linear action of $\mathbf{S}^1$ on $\mathbb{C}\mathrm{P}^1$:

$$[z_0 : z_1] \to [e^{ia\varphi} z_0 : z_1].$$

(S_3): $n = 3$, weights in p_1 and p_2 are $(a, b, -(a + b))$ and $(-a, -b, a + b)$ respectively, where a and b are positive integer, and $\varepsilon_1 = \varepsilon_2$. This action admit $\mathbb{S}^6$ that is almost complex.

Our main result is:

Theorem 1. *Let M be a unitary $2n$–dimensional $\mathbf{S}^1$–manifold with two isolated fixed points. Then the set of weights can only be Z, L_1 or S_3.*

Our proof relies on the fact that $T_{x,y}$ genus is rigid. We consider this method in the next section.

Kosniowski [4] conjectured that a unitary $2n$–dimensional $\mathbf{S}^1$–manifold that is not bound equivariantly has at least $\lceil n/4 \rceil + 1$ isolated fixed points. Theorem

*This research is partially supported by the NSF grant DMS-1400876 and the RFBR grant 15-01-99563.

1 proves the conjecture for $n \leq 4$. Note that this problem was solved only for $\mathbf{T}^n$–manifolds. Namely, Zhi Lü [6] was proved that for torus manifolds there are at least $\lceil n/2 \rceil + 1$ isolated fixed points.

It is not clear, can Kosniowski's conjecture be proved using only the rigidity of the $T_{x,y}$ genus? Perhaps, explicit generators of $U_*^{\mathbf{S}^1}$ from our paper [8] can be useful for that.

2 Rigid Hirzebruch genera

Let U_* be the complex bordism ring with coefficients in $R = \mathbb{Q}$, $\mathbb{R}$, or $\mathbb{C}$. For a closed smooth stably complex manifold X, Hirzebruch [3] defined a multiplicative genus $h(X)$ by a homomorphism $h : U_* \otimes R \to R$.

Recall that according to Milnor and Novikov, two stably complex manifolds are complex cobordant if and only if they have the same Chern numbers. Therefore, for any multiplicative genus h there exists a multiplicative sequence of polynomials $\{K_i(c_1, \ldots, c_i)\}$ such that $h(X) = K_n(c_1, \ldots, c_n)$, where the c_k are the Chern classes of X and $n = \dim_{\mathbb{C}}(X)$ (see [3]).

Let U_*^G be the ring of complex bordisms of manifolds with actions of a compact Lie group G. Then for any homomorphism $h : U_* \otimes R \to R$ we can define the *equivariant genus* h^G, i.e. a homomorphism

$$h^G : U_*^G \otimes R \to K(BG) \otimes R$$

(see details in [5]).

A multiplicative genus h is called *rigid* if for any connected compact group G the equivariant genus $h^G(X) = h(X)$. For the complex case rigidity means that

$$h^G : U_*^G \otimes R \to R \subset K(BG) \otimes R,$$

i.e. $h^G([X, G])$ belongs to the ring of constants. It is well known (see [1, 5]) that $\mathbf{S}^1$-rigidity implies G-rigidity, i.e. it is sufficient to prove rigidity only for the case $G = \mathbf{S}^1$.

For $G = \mathbf{S}^1$, the universal classifying space BG is $\mathbb{C}\mathrm{P}^\infty$, and the ring $K(BG) \otimes R$ is isomorphic to the ring of formal power series $R[[u]]$. Then for any $\mathbf{S}^1-$manifold X and a Hirzebruch genus h we have $h^{S^1}([X, \mathbf{S}^1])$ in $R[[u]]$. In the case of an $\mathbf{S}^1-$action on a unitary manifold X^{2n} with isolated fixed points $p_1 \ldots, p_m$ with weights $w_{i1}, \ldots, w_{in}$, and signs ε_i, $i = 1, \ldots, m$, h^{S^1} can be found explicitly (see [2, 5]):

$$h^{S^1}([X, \mathbf{S}^1]) = S_h(\{w_{ij}\}, u) := \sum_{i=1}^{m} \varepsilon_i \prod_j \frac{H(w_{ij}\, u)}{w_{ij} u}, \tag{1}$$

where H is the characteristic series of h (see [3, 8]). If h is rigid, then from (1) it follows that

$$h(X) = S_h(\{w_{ij}\}, u) \ \text{ for any } \ u. \tag{2}$$

Atiyah and based on the Atiyah-Singer index theorem proved that T_y-genus is rigid [1]. Krichever [5] gives a proof of rigidity of the $T_{x,y}-$genus using global analytic properties of $S_h(\{w_{ij}\},u)$. It is not hard to see that (2) yields the Atiyah-Hirzebruch formula [1, 5, 2] for a unitary $\mathbf{S}^1$-manifold X:

$$T_{x,y}(X)=\sum_{i=1}^{m}\varepsilon_i x^{s_i^+}(-y)^{s_i^-}, \tag{3}$$

where s_i^+, s_i^- are numbers of positive and negative weights $\{w_{ij}\}$.

In [5] is given the following formula for the $T_{x,y}-$genus:

$$H_{x,y}(u)=\frac{u(xe^{u(x+y)}+y)}{e^{u(x+y)}-1}.$$

Let $z:=e^{(x+y)u}$. Then (1), (2), and (3) imply

$$\sum_{i=1}^{m}\varepsilon_i\prod_j\frac{xz^{w_{ij}}+y}{z^{w_{ij}}-1}\equiv\sum_{i=1}^{m}\varepsilon_i x^{s_i^+}(-y)^{s_i^-} \tag{4}$$

3 Proof of the main theorem

Proof. Since we have only two fixed points the graph of weights (see [9]) has two vertices and n edges join these vertices. It implies that $|w_{1i}|=|w_{2i}|$ for all $i=1,\dots,n$. (Actually, it can be derived from (4), where $x=y=1$, see [9, Theorem 1.1].) Moreover, if the set of weights is not Z, then $w_{1i}=-w_{2i}$. Otherwise, $S_{T_{x,y}}(\{w_{ij}\},u)$ cannot be constant. Therefore, for this case formula (4) can be written in the following form:

$$\frac{(xz^{a_1}+y)\cdots(xz^{a_k}+y)(x+yz^{b_1})\cdots(x+yz^{b_\ell})}{(z^{a_1}-1)\cdots(z^{a_k}-1)(z^{b_1}-1)\cdots(z^{b_\ell}-1)}-$$

$$\frac{(x+yz^{a_1})\cdots(x+yz^{a_k})(xz^{b_1}+y)\cdots(xz^{b_\ell}+y)}{(z^{a_1}-1)\cdots(z^{a_k}-1)(z^{b_1}-1)\cdots(z^{b_\ell}-1)}=x^ky^\ell-x^\ell y^k, \tag{5}$$

where $k+\ell=n$ and all a_i and b_j are positive integers.

So for $n=1$ we have L_1. If $n>1$, then $k>0$ and $\ell>0$. For $y=0$ (5) implies

$$a_1+\ldots+a_k=b_1+\ldots+b_\ell \tag{6}$$

Without loss of generality we may assume that $a:=a_1\geq a_i$ for all i and $a_1>b_j$ for all j. (As we mentioned above, we cannot have the equality $a_1=b_j$.) Let $x=-z^a$ and $y=1$. Then (5) yields

$$\frac{(z^{2a}-1)\cdots(z^{a_k+a}-1)(z^a-z^{b_1})\cdots(z^a-z^{b_\ell})}{(z^a-1)\cdots(z^{a_k}-1)(z^{b_1}-1)\cdots(z^{b_\ell}-1)}=(-1)^\ell z^{ak}-(-1)^k z^{a\ell} \tag{7}$$

Since $a>b_j$, we have $a\ell>b_1+\ldots+b_\ell$. Then (7) implies

$$ka=b_1+\ldots+b_\ell \tag{8}$$

Therefore, from (6) we have $a_1 + \ldots + a_k = ka$, $a_1 = \ldots = a_k = a$, k is odd, and ℓ is *even*. Then

$$\frac{(z^a+1)^k(z^{a-b_1}-1)\cdots(z^{a-b_\ell}-1)}{(z^{b_1}-1)\cdots(z^{b_\ell}-1)} = z^{a(\ell-k)}+1 \tag{9}$$

Equation (9) yields $k = 1$, $\ell = 2$ and $a = b_1 + b_2$. Thus, it is the case S_3. □

References

[1] M. F. Atiyah and F. Hirzebruch, Spin manifolds and group actions, in Essays in Topology and Related Subjects, Springer-Verlag, Berlin, 1970, pp. 18-28.

[2] V. M. Buchstaber and N. Ray, The universal equivariant genus and Krichever's formula, *Russian Math. Surveys*, 62:1 (2007), 178 - 180.

[3] F. Hirzebruch, Topological Methods in Algebraic Geometry, 3rd edition, Grundlehren der mathematischen Wissenschaften, no. 131, Springer, Berlin-Heidelberg 1966.

[4] C. Kosniowski, Some formulae and conjectures associated with circle actions, in "Topology Symposium, Siegen 1979" (Proc. Sympos., Univ. Siegen, Siegen, 1979), 331–339, Lecture Notes in Math., 788, Springer, Berlin, 1980.

[5] I. M. Krichever, Formal groups and the Atiyah - Hirzebruch formula, *Izv. Akad. Nauk SSSR Ser. Mat.* **38** (1974), 1289-1304.

[6] Z. Lü, Equivariant cobordism of unitary toric manifolds, in Conference V. Buchstaber–70 "Algebraic topology and Abelian functions", 18–22 June 2013, Moscow, Abstracts, 53–54.

[7] O. R. Musin, On rigid Hirzebruch genera, *Mosc. Math. J.,* **11:1** (2011), 139–147

[8] O. R. Musin, Generators of $\mathbf{S}^1$-bordisms, *Math. USSR Sb.* **44** (1983), 325-334

[9] O.R. Musin, Circle action on homotopy complex projective spaces, *Mathematical Notices,* **28:1** (1980), 533–540.

O. R. Musin, University of Texas Rio Grande Valley, School of Mathematical and Statistical Sciences, One West University Boulevard, Brownsville, TX, 78520.

E-mail address: oleg.musin@utrgv.edu